arXiv:hep-ph/0702062v2 8 Feb 2007

Isospin breaking in scalar and pseudoscalar channels of radiative J/ψ-decays

V.V.Kiselev*
Russian State Research Center “Institute for High Energy Physics”,
Pobeda 1, Protvino, Moscow Region, 142281, Russia
Fax: +7-4967-744937

In the framework of simple assumption on factorizing a mixing of vector state with isoscalar components in effective amplitudes of isospin breaking caused by the electromagnetic quark current, a branching fraction of radiative $J/\psi \to a_0$ transition is evaluated at the level of $3 \cdot 10^{-3}$.

PACS numbers: 13.20.Gd

I. INTRODUCTION

Radiative decays of J/ψ are considered as a source of gluon enhanced channels relevant to searching for glueball states predicted by QCD (see references in [2, 3]). This belief is based on the consideration of diagrams for the annihilation of charm-quarks composing the initial state as shown in Fig. 1a. The two-gluon state can further covert to both gluonic hadrons and light quarks (see Fig. 1b), that produces the mixing of pure glueballs with the ordinary quark matter. A common belief stands that the photon coupling to light quarks causes the suppression of amplitude by α_s due to the additional exchange by the hard gluon (see Fig. 1c), that yields the isoscalar dominance in the radiative J/ψ-decays, since the contribution due to the electromagnetic current of light quarks causing the isospin breaking is suppressed.

However, the exclusive branching fractions, in practice, are actually suppressed with respect to inclusive ones by an order of magnitude, in fact. Therefore, the yields of hadrons with large gluonic components in the radiative decays of vector charmonium could be of the same order as the yield of ordinary hadrons due to diagrams accompanying one in Fig. 1c. This point could be straightforwardly checked by searching for the isospin breaking radiative decays. Indeed, the isospin breaking component of diagrams with electromagnetic current coupled to light quarks (Fig. 1c) does not interfere with the dominant contribution of isoscalar channel (Fig. 1a and b). Therefore, one could test the magnitude of isospin breaking channels by investigating the transitions to isotriplet scalar and pseudoscalar particles[1] (the isospin $T = 1$). The transition to π^0 is measured, and it is suppressed by two orders of magnitude in comparison with decays to isosinglet pseudoscalar particles, which can include large gluonic components. Then, the scalar channel is of interest. It could be represented by the $J/\psi \to a_0\gamma$ mode, for instance. In the present paper we evaluate the branching fraction of this channel in the framework of simple factorizing the isoscalar heavy-light mixing and electromagnetic transition of isoscalar vector light-quark state into the isotriplet scalar. For this purpose in section II we analyze the effective couplings of the isosinglet vector states with isotriplets ones and show that the current data on the radiative decays allow us to extract relevant mixing multiplied by form-factors in order to predict the widths of $J/\psi \to a_0\gamma$. The result derived is discussed in section III.

II. EVALUATION

Let us introduce effective dimensionless couplings $\widetilde{F}$ and F of vector isoscalar particle $V = \omega$, ϕ, J/ψ with the neutral isotriplet pseudoscalar π^0 and scalar a_0 in amplitudes of radiative decays as follows:

$$\widetilde{\mathcal{M}}(V \to \pi^0\gamma) = \frac{\sqrt{4\pi\alpha}}{2m_V}\,\widetilde{F}_{V\pi^0}\,\widetilde{\mathcal{F}}_{\mu\nu}\mathcal{V}^{\mu\nu}, \tag{1}$$

$$\mathcal{M}(V \to a_0\gamma) = \frac{\sqrt{4\pi\alpha}}{2m_V}\,F_{Va_0}\,\mathcal{F}_{\mu\nu}\mathcal{V}^{\mu\nu}, \tag{2}$$

where $\mathcal{F}_{\mu\nu}$ is the strength tensor of electromagnetic field $\mathcal{A}_\mu$

$$\mathcal{F}_{\mu\nu} = \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu,$$

$\widetilde{\mathcal{F}}_{\mu\nu}$ denotes its dual tensor

$$\widetilde{\mathcal{F}}_{\mu\nu} = \frac{1}{2}\,\epsilon_{\mu\nu\alpha\beta}\mathcal{F}^{\alpha\beta},$$

while $\mathcal{V}^{\mu\nu}$ is the strength tensor of vector $\mathcal{V}_\mu$. For brevity we include the product of effective electric charges Q_V and Q_{π^0,a_0} into the definition of couplings $\widetilde{F}$ and F. Then, we easily get the widths of decays

$$\Gamma(V \to \pi^0\gamma) = \frac{\alpha}{3}\,\frac{k_0^3}{m_V^2}\,\widetilde{F}^2_{V\pi^0}, \tag{3}$$

$$\Gamma(V \to a_0\gamma) = \frac{\alpha}{3}\,\frac{k_0^3}{m_V^2}\,F^2_{Va^0}, \tag{4}$$

with k_0 being the photon energy

$$k_0 = \frac{m_V^2 - m_{\pi^0,a_0}^2}{2m_V}.$$

---

*Electronic address: Valery.Kiselev@ihep.ru

[1] Similar mechanism contributes to hadronic decays of J/ψ and ψ', if one considers the virtual photons as it was studied in [1], wherein a role of isospin breaking was also emphasized.



FIG. 1: The annihilation in radiative decays of J/ψ: a) the gluonic channel, b) the gluon conversion into light quarks in the isoscalar channel, c) the gluon conversion into light quarks with breaking off the isospin.

The experimental values of measured branching fractions [4] approximately yield the effective couplings to the neutral pion[2]

$$\begin{aligned} \widetilde{F}_{\omega\pi^0} &= 1.86 \pm 0.03, & (5)\\ \widetilde{F}_{\phi\pi^0} &= 0.135 \pm 0.004, & (6)\\ \widetilde{F}_{\psi\pi^0} &= (1.81 \pm 0.36)\, 10^{-3}, & (7) \end{aligned}$$

that spectacularly shows order by order decreasing of effective couplings with stepping to more heavy vector state. Such the regularity can be understood in a simple manner.

The diagram shown in Fig. 2 describes the mixing of vector isoscalar quark states in the lowest order of QCD coupling constant. Therefore, the mixing does exist, though its evaluation is beyond the perturbation theory. The hierarchy of couplings in (5)–(7) implies that we could represent the transition of more heavy states ϕ and J/ψ by the mixing with ω and further radiative decay, i.e.

$$\widetilde{F}_{V\pi^0} = \mathcal{K}_{V\omega}\, \widetilde{F}_{\omega\pi^0}. \qquad (8)$$

The $\mathcal{K}$-factors include both the mixing amplitude and form-factors caused by the virtuality of ω-propagation. Therefore, these factors cannot be straightforwardly extracted from the phenomenology of ω–ϕ-mixing, say.



FIG. 2: The mixing of vector isoscalar quark states in the lowest order of QCD coupling constant.

In this way we find the $\mathcal{K}$-factors

$$\mathcal{K}_{\phi\omega} = (7.2 \pm 0.2)\, 10^{-2}, \quad \mathcal{K}_{\psi\omega} = (10 \pm 2)\, 10^{-4}. \qquad (9)$$

Supposing the same mechanism for the radiative decay to isotriplet scalar meson a_0, we put

$$F_{Va_0} = \mathcal{K}_{V\omega}\, F_{\omega a_0}, \qquad (10)$$

while the experimental value of $\Gamma(\phi \to a_0\gamma)$ yields

$$F_{\phi a_0} = 1.5 \pm 0.1, \qquad (11)$$

that results in effective value of coupling for the kinematically forbidden decay

$$F_{\omega a_0} = 21 \pm 2, \qquad (12)$$

as well as the quantity of interest

$$F_{\psi a_0} = (20 \pm 5)\, 10^{-3}, \qquad (13)$$

giving

$$\mathcal{B}r(J/\psi \to a_0\gamma) = (3.1 \pm 1.5)\, 10^{-3}, \qquad (14)$$

where the uncertainty is dominated by experimental one in the measurement of radiative decays of J/ψ to pion as well as ϕ to a_0, while the theoretical uncertainty could be additionally included *ad hoc* at the level of 20%, characteristic for the exclusive modelling the virtual process of mixing and form-factors.

III. DISCUSSION AND CONCLUSION

The only experimental data available to the moment for the relevant process is presented by

$$\mathcal{B}r(J/\psi \to \gamma\pi^+\pi^-2\pi^0) = (8.3 \pm 3.1)\, 10^{-3}, \qquad (15)$$

that should be compared with expected

$$\begin{aligned} &\mathcal{B}r(J/\psi \to \gamma a_0) \times \mathcal{B}r(a_0 \to \eta\pi^0)\times \\ &\qquad \mathcal{B}r(\eta \to \pi^+\pi^-\pi^0) \;=\; (0.7 \pm 0.4)\, 10^{-3}, \end{aligned}$$

---

[2] The values in (5)–(7) reproduce those of derived in [1] after appropriate rescaling in definition: $g = \widetilde{F}\,\sqrt{4\pi\alpha}$.

while the neutral modes of η decay in such final states of J/ψ decay are not available yet. We see that the resonant a_0-contribution due to the cascade decay with the $\gamma\pi^+\pi^-2\pi^0$ final state could give about 10% of branching ratio summed over all modes as we expect.

The main conclusion of estimate obtained is the following: the ordinary quark states could compose a considerable fraction of radiative J/ψ-decay in contrast to usual assumption on the enhancement of gluonic component. Moreover, the isospin breaking effects could be essential. Thus, the simple treatment of resonances in the scalar channel as the probable glueball states is under question.

The critical point of estimate is the factorization of mixing form-factors and effective couplings. This suggestion is natural. The resulting mixing factors are small, that should support the assumption. However, the suppression of mixing could lead to the method suffers from small effects breaking the factorization. For instance, the light quark states, ω, a_0 or ϕ with the hidden strangeness could get a small admixture of four-quark states. Then, the admixture would break the factorization. Nevertheless, we expect that the four-quark mechanism gets a significant suppression, since in the exclusive modes one should include the annihilation of additional quark-antiquark pair. Therefore, we expect that the method used is able to present the real estimate for the isospin breaking effect in the radiative J/ψ-decay to a_0.

The author is grateful to prof. A.M. Zaitsev for initiating the consideration of problem and useful discussions.

This work is partially supported by the Russian Foundation for Basic Research, grant 07-02-00417-a.

---

[1] Q. Zhao, G. Li and C. H. Chang, Phys. Lett. B **645**, 173 (2007) [arXiv:hep-ph/0610223].

[2] F. E. Close, Rept. Prog. Phys. **51**, 833 (1988);
C. Amsler and F. E. Close, Phys. Lett. B **353**, 385 (1995) [arXiv:hep-ph/9505219];
C. Amsler and F. E. Close, Phys. Rev. D **66**, 010001 (2002) arXiv:hep-ph/9507326;
F. E. Close, G. R. Farrar and Z. P. Li, Phys. Rev. D **66**, 010001 (2002) arXiv:hep-ph/9610280;
F. E. Close and A. Kirk, Phys. Lett. B **483**, 345 (2000) [arXiv:hep-ph/0004241];
F. E. Close and N. A. Tornqvist, J. Phys. G **28**, R249 (2002) [arXiv:hep-ph/0204205].

[3] V. V. Anisovich and A. V. Sarantsev, Phys. Rev. D **66**, 010001 (2002) arXiv:hep-ph/0204328.

[4] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33**, 1232 (2006).